

PUBLICATIONS

1996

FIELD DAY REPORT - 1996

TEXAS A&M UNIVERSITY AGRICULTURAL RESEARCH and EXTENSION CENTER at OVERTON

**Texas Agricultural Experiment Station
Texas Agricultural Extension Service**

Overton, Texas

April 18, 1996

Research Center Technical Report 96-1

All programs and information of the Texas Agricultural Experiment Station and Texas Agricultural Extension Service are available to everyone without regard to race, color, religion, sex, age, or national origin.

Mention of trademark or a proprietary product does not constitute a guarantee or a warranty of the product by the Texas Agricultural experiment Station or Texas Agricultural Extension Service and does not imply its approval to the exclusion of other products that also may be suitable.

SEASONAL FORAGE PRODUCTION OF SOD-SEEDED CLOVER MIXTURES

G. R. Smith, I. J. Pemberton, and J. Gilbert

Background. Annual clovers are useful components of east Texas pasture systems but the productive growing season for an individual clover species is often short. Crimson clover produces forage early, usually in the months of March and April. Arrowleaf clover has a longer and later production season than crimson but does not grow rapidly in the early spring. Red and white clover are perennials that generally act like annuals under east Texas climate and soil conditions. Both red and white clover are most productive from late April to early June in east Texas. Ball clover grows rapidly in April and May but usually produces little forage before April. Mixtures of two or more different clovers can be used to extend the forage legume growing season. The objective of these experiments was to determine the seasonal distribution of forage yield from mixtures of two clovers.

Research Findings. 'Kenland' red clover, 'Dixie' crimson, 'Yuchi' arrowleaf, 'Regal' white, common ball clover, and sixteen clover mixtures were drilled into a 'Coastal' bermudagrass sod on 22 Oct. 1992, on 15 Oct. 1993, and 14 Oct. 1994. Treatments were arranged in a randomized complete block design with four replications. Fertilizer was applied prior to planting according to soil test recommendations (400 lbs/ac 0-20-20 and 1 lb/ac boron in 1992; 400 lbs/ac 0-20-33 and 1 lb/ac boron in 1993 and 1994). A small-plot drill with six double disk openers spaced 9 inches apart was used to place the clover seed 0.5 inch deep. Seed were inoculated with specific *Rhizobium* inoculant prior to planting. Plots containing only single clovers were planted at 14, 4, 20, 14, and 6 lbs of seed per acre for arrowleaf, ball, crimson, red, and white clover, respectively. All mixtures were planted with crimson clover at either 50% or 75% of the full seeding rate.

Plots were harvested at about 4-week intervals beginning in March with a rotary mower set to cut 2.5 inches above ground level. Fresh weight of forage from each plot was recorded at each harvest date. Subsamples were also weighed at each harvest, dried at 140 F for 48 hours, then weighed again to determine percent dry matter. Dry forage yield per acre was calculated.

Dixie crimson alone or in mixtures produced the most forage at the first harvest in all years (1995 data shown in Fig. 1, others years not shown). Red and white clover were consistently low yielding at the first harvest compared to crimson or mixtures containing crimson. The mixture of crimson and Kenland red clover was very productive in all years with good distributions of forage yield over the growing season. The D50/K50 mixture (10 lbs/ac Dixie

crimson + 7 lbs/ac Kenland red clover) was generally as productive as the D75/K75 mixture and seed costs were \$21.65 per acre for the D50/K50 mixture compared to \$32.47 per acre for the D75/K75 mixture.

Mixtures of crimson with Regal white clover, Yuchi arrowleaf, or common ball clover generally resulted in lower yields at the last two harvests and lower total yields compared to the red clover mixtures. Forage production of Yuchi arrowleaf and mixtures of Yuchi and crimson clover were reduced due to fungal and virus diseases of arrowleaf clover.

Application. Mixtures of two or more clovers can result in more high quality forage available to grazing animals over a longer period of time. The best clover mixture evaluated from 1993-95 was Dixie crimson (10 lbs/ac) plus Kenland red clover (7 lbs/ac). In this mix, crimson clover was productive by early March and red clover extended the clover grazing season into late May or early June.

Fig. 1. Forage production of clover mixtures sod-seeded in Coastal bermudagrass. Letter number combinations indicate clover species and percent of full seeding rate used. Data from 1995.

