

PUBLICATIONS

1986

FORAGE AND LIVESTOCK RESEARCH - 1986

RESEARCH CENTER TECHNICAL REPORT 86-1

Texas A&M University Agricultural Research
and Extension Center at Overton

Texas Agricultural Experiment Station
Texas Agricultural Extension Service

Overton, Texas

April 24, 1986

All programs and information of the Texas Agricultural Experiment Station and Texas Agricultural Extension Service are available to everyone without regard to race, color, religion, sex, age, or national origin.

Mention of trademark or a proprietary product does not constitute a guarantee or a warranty of the product by the Texas Agricultural Experiment Station or Texas Agricultural Extension Service and does not imply its approval to the exclusion of other products that also may be suitable.

RESPONSE OF COASTAL BERMUDAGRASS TO A HIGH SULFUR CONTENT
SULPHATE OF POTASH-MAGNESIA

V. A. Haby, J. V. Davis, and A. T. Leonard

SUMMARY

Coastal bermudagrass response to a sulfur-magnesium treatment combination was evaluated on a Gallime fine sandy loam soil containing 8.3 and 24.2 ppm sulfur (S) and magnesium (Mg), respectively, in the 0-6 inch depth. Rates of 0, 40, 80, and 120 lb S/ac and 0, 2.95, 5.89, and 8.8 lb Mg/ac were applied to the same plots as a 57% S content sulphate of potash-magnesia. Nitrogen (N), phosphorus, and potassium were equalized over all plots initially at rates of 100, 100, and 250 lb of N, P_2O_5 , and K_2O /ac, respectively, and N at the rate of 100 lb/ac was applied after each cutting. Three harvests of grass were made. Yields averaged 5004, 2715, and 5237 lb/ac at each cutting, respectively, and totalled an average of 12,950 lb/ac for the three cuttings. The S-Mg combination treatment had no statistically significant effect on yield of grass.

OBJECTIVE

Evaluate the response of Coastal bermudagrass to a combination of sulfur and magnesium on a Gallime fine sandy loam soil.

PROCEDURE

A Gallime fine sandy loam soil having a surface pH of 5.1 was selected for this study. This soil contained 8.3 and 24.2 ppm sulfur (S) and magnesium (Mg), respectively, in the surface six inch depth. Rates of S were 0, 40, 80, and 120 lb/ac and were broadcast onto the surface of 10 x 20 foot plots. Magnesium levels were not equalized over all plots, but were allowed to increase as the S rate increased. Sulphate of potash magnesia containing 8.4% K_2O , 4.2% Mg, 8.4% sulfate-S and a total of 57% S was the fertilizer material used to vary the sulfur rate. Potassium treatment was equalized on all plots at 250 lb K_2O /ac. Nitrogen and phosphorus (P_2O_5) were each applied at 100 lb/ac, initially. Nitrogen at 100 lb/ac was applied after each of the first two cuttings. Yield samples were taken in July, August, and November.

RESULTS

Responses of Coastal bermudagrass to the sulfur-magnesium rates are indicated in Table 1. Yields by treatment were not significantly different. This indicated that available S and Mg were adequately supplied in the soil, if not in the top 6 inches, possibly in the lower depths. Sulfur and Mg in the harvested grass have not been analyzed.

TABLE 1. EFFECT OF SULFUR-MAGNESIUM COMBINATION RATES ON YIELD OF COASTAL BERMUDAGRASS

Treatment S - Mg	Yield of Coastal bermudagrass dry matter			
	Harvest 1	Harvest 2	Harvest 3	Total
	-----lb/ac-----			
0 - 0	4860	2517	5161	12530
40 - 2.95	4590	2835	5079	12500
80 - 5.89	5217	2638	5614	13460
120 - 8.80	5348	2871	5094	13310
Mean	5004	2715	5237	12950
L.S.D. (p=0.10)	N.S.	N.S.	N.S.	N.S.
C.V. %	12.7	13.7	8.8	6.2