




PUBLICATIONS

1983



Forage Research in Texas

1983

Nutritive Value of White Clover Varieties

F. M. Rouquette, Jr., G. R. Smith, C. Gilbert, and R. Gillespie

SUMMARY

Sixteen white clover varieties were evaluated for percent protein and in vitro digestible dry matter (IVDDM) during the spring-summer growing period. Protein ranged from about 30% in March to about 20% in mid-July. IVDDM declined slowly from about 80% in March to approximately 70% in mid-July.

Introduction

White clovers are grown in the eastern one-third of Texas on bottomland-type soils. Although the primary objective associated with white clover forage production is that of summer persistence and survival, the nutritive value of this clover as influenced by season is also useful information to the livestock producer. The primary objective of this trial was to monitor percent protein and IVDDM of selected white clover varieties throughout the growing season.

Procedure

Seventeen white clover varieties were seeded at the Texas A&M University Agricultural Research and Extension Center in a randomized complete block design with four replications. Plots were broadcast seeded at the rate of six lbs/acre in mid-October on a bottomland site containing a Thenas fine sandy loam soil. Inoculum (Type B) was applied to the seed pre-plant at 3X the recommended rate. Lime was applied at 1.5 tons/acre and plots were fertilized at planting with 24-96-96 lbs/acre of $N-P_2O_5-K_2O$, respectively. Plots were clipped to a 2.25 inch stubble with a rotary mower. Dry matter yields were determined and subsamples were taken for chemical analyses. Clover samples were analyzed for protein via micro Kjeldahl and IVDDM via modified Barnes technique.

Results and Discussion

Table 1 shows the monthly rainfall during the initial establishment year (79-80) and the data collection year (80-81). Although the total rainfall during 80-81 was less than the 13-year average, the precipitation that occurred during the May-July months was above normal, and primarily responsible for the extended forage production season. Forage dry matter harvested at monthly intervals is presented in Table 2 (1). Forage produced during this growing season was the result of clover survival during the previous year. Dry matter production during the year of establishment was about one-half that shown in Table 2.

Table 3 shows the influence of season on percent protein of the white clover varieties. Forage harvested on March 20 had more than 30% protein. Substantial decreases in percent protein occurred at the May 21 harvest and again at the July 16 harvest. However, since the absolute protein values remained at nearly 20%, this protein reduction due to chronological and physiological maturity would not be expected to detrimentally affect animal performance.

IVDDM of the clover varieties is presented in Table 4. The decline in percent IVDDM followed a similar trend to that of percent protein. Percent IVDDM ranged from slightly more than 80% on March 20 to about 70% on July 16. Both Fla. XPL and S.C. Nematode Tolerant had IVDDM values of approximately 80% on July 16; whereas, most of the other varieties had IVDDM values of about 70%. The May 21 harvest date produced most of the low IVDDM values during the measurement period. This may have been due to the relatively small amount of rainfall which occurred during the previous month; or it may have been due to advancing physiological maturity as evidenced by an increasing amount of flowering which occurred in early to mid-May. The quality of these forages, as represented by both protein and IVDDM, remained relatively high throughout the trial period and individual animal performance should be greatly improved over that performance obtained from forages such as warm-season perennial grasses.

Literature Cited

1. Smith, G. R., F. M. Rouquette, Jr., and E. R. Shipe. 1982. Forage production of red and white clover in East Texas.

Table 1. Precipitation by months at Overton, Texas for a two year period.

Months	Years		13 year Avg.
	1979-80	1980-81	
-----inches-----			
September	6.6	3.2	4.7
October	2.4	1.9	3.7
November	4.3	3.5	3.8
December	5.1	1.5	3.3
January	3.6	1.0	4.0
February	2.6	2.8	3.3
March	2.6	2.8	4.1
April	4.3	2.0	4.7
May	6.1	7.8	3.9
June	2.1	4.7	4.0
July	1.3	5.4	2.9
August	0.9	0.7	1.9

Table 2. White Clovers Grown on a Prepared Seedbed.

Entry	Harvest Date					Total
	3-20	4-13	5-21	6-12	7-16	
-----lbs DM/A-----						
Fla. XPL ²	1343	1610	1412	1480	1329	7174 a ¹
L8-100 ²	1571	1457	1010	1408	1597	7043 ab
Lucky	1583	1418	1109	1284	1613	7007 ab
L8-101 ²	1511	1516	1084	1437	1435	6983 ab
Calif. Ladino	1499	1501	1171	1309	1451	6931 ab
S.C. Med. ²	1391	1424	1061	1479	1541	6896 abc
S.C. Nem. Tol. ²	1535	1497	1133	1327	1386	6878 abc
Arcadia	1343	1427	1152	1367	1526	6815 abcd
Tillman	1499	1354	950	1187	1404	6394 bcde
K6-8 Ladino ²	1367	1224	989	1192	1521	6293 cdef
Regal	1223	1295	1022	1235	1468	6243 def
Sacramento	1139	1302	1179	1161	1275	6056 ef
La. S-1	1355	1646	1477	1265	0	5743 fg
Nolin's Imp.	1151	1678	1425	1206	0	5460 gh
Tamar	1103	1572	1158	1133	0	4966 h
Rector	1427	1282	1097	1041	0	4847 h

C.V. = 62.2%

¹ Numbers within a column not followed by the same letter differ (P<.05) according to the Duncan's Multiple Range test.

² Experimental varieties or germplasm.

Table 3. Percent protein of selected white clovers during the growing season.

Entry	Harvest Date				
	3-20	4-13	5-21	6-12	7-16
Fla. XPL	32.8	30.1	25.0	23.6	20.0
L8-100	31.3	33.7	25.8	25.8	18.8
Lucky	32.5	29.4	26.8	24.1	21.2
L8-101	34.8	28.9	28.4	25.8	21.5
Calif. Ladino	32.5	33.0	26.6	23.6	19.3
S.C. Med.	33.6	32.6	27.6	24.3	20.6
S.C. Nem. Tol.	31.6	32.5	29.0	24.6	21.1
Arcadia	31.5	34.5	25.9	23.3	18.8
Tillman	32.3	33.7	28.8	25.0	23.1
K6-8 Ladino	32.2	33.2	26.8	23.6	20.7
Regal	34.9	30.0	25.3	24.9	21.7
Sacramento	33.0	28.5	25.1	24.2	19.4
La. S-1	30.4	31.2	22.4	22.6	-
Nolin's Imp.	32.6	29.5	24.0	21.1	-
Tamar	32.4	30.5	23.3	21.7	-
Rector	34.3	32.6	-	20.4	-

Table 4. Percent in vitro digestible dry matter of selected white clovers during the growing season.

Entry	Harvest Date				
	3-20	4-13	5-21	6-12	7-16
Fla. XPL	83.3	74.4	71.8	69.8	80.7
L8-100	80.0	74.3	71.0	77.6	75.9
Lucky	81.4	76.6	75.4	76.5	73.4
L8-101	82.3	74.0	76.8	76.2	74.4
Calif. Ladino	84.5	75.3	65.3	75.4	67.9
S.C. Med.	82.9	72.8	70.8	73.6	67.1
S.C. Nem. Tol.	80.4	72.0	71.0	77.1	79.4
Arcadia	80.4	78.6	66.1	72.9	72.6
Tillman	81.7	76.5	72.0	75.9	68.8
K6-8 Ladino	82.3	73.7	68.1	74.9	72.6
Regal	79.5	74.3	68.6	74.5	72.8
Sacramento	81.9	73.0	66.6	71.0	74.6
La. S-1	78.1	73.8	58.8	68.4	-
Nolin's Imp.	81.8	70.8	65.1	74.6	-
Tamar	81.6	68.1	66.9	69.3	-
Rector	79.4	73.5	-	66.8	-