

PUBLICATIONS

1987

FRUIT AND NUT CROPS RESEARCH IN TEXAS, 1987

Page	Participating Scientists	Crops
3, 5	David H. Byrne	Peach
3, 5	Terry Bacon	Plums
7	J. Dan Hanna	Apricots
9	Calvin G. Lyons	Grapes
11, 12	T. Lynn Littleton	Pecans
10	G. R. McEachern	
19, 20, 48	Bert Johnson	
12	J. Benton Storey	
48	Berry Tompkins	
15	R. D. Marquard	Pecan
17	L. Austin Stockton	Grapes Apples
19, 20, 21, 23	John A. Lipe	Peach
19, 20	Dusty Menzies	Pecan

COMPILED AND EDITED BY:

Robert E. Rouse
 Texas Agricultural Experiment Station
 2415 East Highway 83
 Weslaco, TX 78596

David H. Byrne
 Department of Horticulture
 Texas A&M University
 College Station, TX 77843

The Texas Agricultural Experiment Station, Neville P. Clarke, Director,
 Texas A&M University System, College Station, TX.

SUBJECT TOPIC: Stonefruit Rootstock Development

INVESTIGATOR(S): David H. Byrne - TAES, College Station
 Terry Bacon - TAES, College Station
 T. Glynn Littleton - TAES, College Station

CROP(S):

1. Peaches
2. Nectarines
3. Plums
4. Apricots

ABSTRACT:

Objectives:

1. Identify useful rootstock materials tolerant to various edaphic and pathogenic stresses.
2. Collect native wild Prunus species to evaluate as possible sources of tolerance to the edaphic/pathogenic stresses encountered in Texas.

General Approach:

1. The development of screening procedures and the assessment of the variability and the level of tolerance to various nutritional stresses have been emphasized.
2. The collection and establishment of native Prunus accessions from throughout Texas and from other habitats is the first step. Once established, the accessions will be evaluated for tolerances and horticultural traits (vigor, ease of propagation, suckering, etc.)

Findings:

1. Preliminary work was shown variability in Prunus for tolerance to alkalinity, salinity and acidity (Al and Mn toxicity). A laboratory procedure to screen for tolerance to alkalinity is being tested. Field screening plots are being developed. Nemared seedlings when subjected to moderate alkalinity stress showed a 60% decrease in dry weight after 31 weeks. Under the same conditions, hybrid ('Titan' x Nemaguard) seedlings showed little chlorosis and no growth reduction. The hybrid was more variable, more vigorous, and produced fewer suckers than Nemared or Nemaguard.
2. Accessions of Prunus angustifolia, P. umbellata, P. mexicana, P. rivularis, P. texana and P. minutiflora have been collected from various parts of the state. More accessions are being collected especially from areas with alkaline soils, severe soil-borne disease

problems and with periodic waterlogged conditions. Once established these will be evaluated for tolerances to various soil stresses and horticultural characteristics.

INVESTIGATOR(S): David H. Byrns - TACS, College Station
Terry Bacon - TACS, College Station
T. Lynn Listeron - TACS, College Station

- 1. Peaches
- 2. Nectarines
- 3. Plums
- 4. Apricots

ABSTRACT:

Objectives:

- 1. Identify useful rootstock materials tolerant to various edaphic and pathogenic stresses.
- 2. Collect native wild *Prunus* species to evaluate as possible sources of tolerance to the edaphic/pathogenic stresses encountered in Texas.

General Approach:

- 1. The development of screening procedures and the assessment of the variability and the level of tolerance to various nutritional stresses have been emphasized.
- 2. The collection and establishment of native *Prunus* accessions from throughout Texas and from other habitats is the first step. Once established, the accessions will be evaluated for tolerances and horticultural traits (vigor, ease of propagation, suckering, etc.).

Findings:

- 1. Preliminary work was shown variability in *Prunus* for tolerance to alkalinity, salinity and acidity (AI and MI toxicity). A laboratory procedure to screen for tolerance to alkalinity is being tested. Field screening plots are being developed. Nemared seedlings when subjected to moderate alkalinity stress showed a 60% decrease in dry weight after 31 weeks. Under the same conditions, hybrid ('Titan' x Nemared) seedlings showed little chlorosis and no growth reduction. The hybrid was more variable, more vigorous, and produced fewer suckers than Nemared or Nemared.
- 2. Accessions of *Prunus angustifolia*, *P. umbellata*, *P. mexicana*, *P. tinctoria*, *P. texana* and *P. minorifolia* have been collected from various parts of the state. More accessions are being collected especially from areas with alkaline soils, severe soil-borne disease